

## Epidermal Langerhans Cell Migration in CD44-Deficient Mice

isodisomy of chromosome 17 and the first example of UPD underlying JEB-PA.

## CONFLICT OF INTEREST

The authors state no conflict of interest.

## ACKNOWLEDGMENTS

We thank Professor Jouni Uitto for his critical comments on the present case, Ms Yuko Hayakawa and Ms Yuki Miyamura for their technical assistance, and Mr Michael O'Connell for his proofreading. This work was supported in part by grants-in-aid from the Ministry of Education, Culture, Sports, Science and Technology to H. Shimizu (Kiban A 21249063).

**Ken Natsuga<sup>1</sup>, Wataru Nishie<sup>1</sup>, Ken Arita<sup>1</sup>, Satoru Shinkuma<sup>1</sup>, Hideki Nakamura<sup>1</sup>, Shogo Kubota<sup>2</sup>, Sumihisa Imakado<sup>3</sup>, Masashi Akiyama<sup>1</sup> and Hiroshi Shimizu<sup>1</sup>**

<sup>1</sup>Department of Dermatology, Hokkaido University Graduate School of Medicine, Sapporo, Japan; <sup>2</sup>Department of Pediatrics, Japanese Red Cross Medical Center, Tokyo, Japan and <sup>3</sup>Department of Dermatology, Japanese Red Cross Medical Center, Tokyo, Japan  
E-mail: [natsuga@med.hokudai.ac.jp](mailto:natsuga@med.hokudai.ac.jp)

## REFERENCES

- Castori M, Floriddia G, Pisaneschi E *et al.* (2008) Complete maternal isodisomy causing reduction to homozygosity for a novel LAMB3 mutation in Herlitz junctional epidermolysis bullosa. *J Dermatol Sci* 51:58–61
- Fassihi H, Lu L, Wessagowit V *et al.* (2006) Complete maternal isodisomy of chromosome 3 in a child with recessive dystrophic epidermolysis bullosa but no other phenotypic abnormalities. *J Invest Dermatol* 126: 2039–43
- Fassihi H, Wessagowit V, Ashton GH *et al.* (2005) Complete paternal uniparental isodisomy of chromosome 1 resulting in Herlitz junctional epidermolysis bullosa. *Clin Exp Dermatol* 30:71–4
- Fine JD, Eady RA, Bauer EA *et al.* (2008) The classification of inherited epidermolysis bullosa (EB): Report of the Third International Consensus Meeting on Diagnosis and Classification of EB. *J Am Acad Dermatol* 58:931–50
- Genuardi M, Tozzi C, Pomponi MG *et al.* (1999) Mosaic trisomy 17 in amniocytes: phenotypic outcome, tissue distribution, and uniparental disomy studies. *Eur J Hum Genet* 7:421–6
- Groves RW, Liu L, Dopping-Hepenstal PJ *et al.* (2010) A homozygous nonsense mutation within the dystonin gene coding for the coiled-coil domain of the epithelial isoform of BPAG1 underlies a new subtype of autosomal recessive epidermolysis bullosa simplex. *J Invest Dermatol* 130:1551–7
- Iacovacci S, Cicuzza S, Odorisio T *et al.* (2003) Novel and recurrent mutations in the integrin beta 4 subunit gene causing lethal junctional epidermolysis bullosa with pyloric atresia. *Exp Dermatol* 12:716–20
- Kotzot D, Utermann G (2005) Uniparental disomy (UPD) other than 15: phenotypes and bibliography updated. *Am J Med Genet A* 136:287–305
- Lebre AS, Moriniere V, Dunand O *et al.* (2009) Maternal uniparental heterodisomy of chromosome 17 in a patient with nephropathic cystinosis. *Eur J Hum Genet* 17: 1019–23
- Nakamura H, Sawamura D, Goto M *et al.* (2005) Epidermolysis bullosa simplex associated with pyloric atresia is a novel clinical subtype caused by mutations in the plectin gene (PLEC1). *J Mol Diagn* 7:28–35
- Pulkkinen L, Bullrich F, Czarnecki P *et al.* (1997) Maternal uniparental disomy of chromosome 1 with reduction to homozygosity of the LAMB3 locus in a patient with Herlitz junctional epidermolysis bullosa. *Am J Hum Genet* 61:611–9
- Rio M, Ozilou C, Cormier-Daire V *et al.* (2001) Partial maternal heterodisomy of chromosome 17q25 in a case of severe mental retardation. *Hum Genet* 108:511–5
- Takizawa Y, Pulkkinen L, Chao SC *et al.* (2000) Mutation report: complete paternal uniparental isodisomy of chromosome 1: a novel mechanism for Herlitz junctional epidermolysis bullosa. *J Invest Dermatol* 115: 307–11
- Takizawa Y, Pulkkinen L, Shimizu H *et al.* (1998) Maternal uniparental meroisodisomy in the LAMB3 region of chromosome 1 results in lethal junctional epidermolysis bullosa. *J Invest Dermatol* 110:828–31
- Varki R, Sadowski S, Pfendner E *et al.* (2006) Epidermolysis bullosa. I. Molecular genetics of the junctional and hemidesmosomal variants. *J Med Genet* 43:641–52

# CD44-Deficient Mice Do Not Exhibit Impairment of Epidermal Langerhans Cell Migration to Lymph Nodes after Epicutaneous Sensitization with Protein

*Journal of Investigative Dermatology* (2010) **130**, 2674–2677; doi:10.1038/jid.2010.170; published online 24 June 2010

## TO THE EDITOR


CD44 is a type I transmembrane protein that binds extracellular matrix nonsulfated glycosaminoglycan hyaluronan and has an important role in cell adhesion and migration (Isacke, 2002). Thus, CD44 is involved with leukocyte egress, tumor invasiveness, and metastasis (Isacke, 2002).

The role of CD44 in epidermal Langerhans cell (LC) migration to drain-

ing lymph nodes (LNs) was first evaluated by an antibody blocking system. Antibodies against CD44 epitopes inhibited emigration of LCs from the epidermis and prevented cultured LC binding to T-cell zones in LN-frozen sections (Weiss *et al.*, 1997). In a CD44-deficient mouse system, CD44 deficiency did not impair LC emigration from the epidermis, but significantly influenced their LN homing (Mummert

*et al.*, 2004). In recent years, there has been significant progress in understanding the characteristics and kinetics of LCs. It is known that there are two kinds of Langerin<sup>+</sup> dendritic cells (DCs) (definition of LCs): one resides in the epidermis and another resides in the dermis (Bursch *et al.*, 2007). They show different migration patterns to draining LNs after immunization. Dermal Langerin<sup>+</sup> DC migration peaks early at 24 hours, whereas peak migration of epidermal LC is delayed until

Abbreviations: DC, dendritic cell; LC, Langerhans cell; LN, lymph node; Th, T helper




**Figure 1. Numbers of dendritic cells (DCs) in draining lymph nodes (LNs).** (a) CD44 staining of peripheral blood leucocytes of  $CD44^{+/+}$  (dash),  $CD44^{P256/+}$  (dotted), and  $CD44^{P256/P256}$  (solid) mice, and unstained control (shaded) are shown. (b) All CD44 exons were subjected to DNA sequencing, and a T-to-A transversion mutation within exon 2 was found (asterisk denotes premature termination). (c) Reverse transcriptase-PCR amplification between exon 1 and exon 5. (d, e, g) Draining LNs were isolated from naive mice (d) or 96 hours after immunization (e, g).  $CD11c^{+}$  cells were gated. Results are expressed as numbers per six draining LNs per mouse. The pooled data of three to five experiments are shown. (f) LNs were isolated from  $CD44$ -deficient mice 96 hours after patch application of ovalbumin (OVA) (dash) or OVA-DQ (solid). Langerin $^{+}$  cells were gated. (g) Langerin $^{high} CD103^{-}$  DCs were gated.

4 days after immunization (Shklovskaya *et al.*, 2008). As previous reports failed to define LCs by specific markers and analyzed LNs for no more than 3 days, we sought to re-evaluate the migration of epidermal LCs in  $CD44$ -deficient mice using a well-established protein-patch model (Chen *et al.*, 2009). Protein antigens rather than haptens were chosen because protein antigens will cross the keratin barrier much slower, owing to their larger size and hydrophilic character, thus giving a bigger chance for epidermal LCs to capture and present them. All animal experiments were approved by the Animal Care Committee of the Medical College of National Taiwan

University. Both  $CD44$ -deficient (P256) and wild-type C57BL/6 mice were from the National Mouse Mutagenesis Program Care Facility of Academia Sinica. A T to A transversion mutation within exon 2 caused a premature termination that resulted in a severely truncated 72-amino-acid  $CD44$  protein, instead of the 778-residue wild-type  $CD44$  protein (Figure 1a and b). Trans-splicing of mRNA between exon 1 and other parts of the standard region of  $CD44$ , which will escape the premature termination in exon 2, did not occur in P256 mice as monitored by reverse transcriptase-PCR amplification between exon 1 and exon 5 (Figure 1c).

First, the numbers of Langerin $^{+}$  DCs in draining LNs of naive  $CD44$ -deficient and wild-type C57BL/6 mice were examined. The numbers of  $CD8^{+}$  DCs and total  $CD11c^{+}$  DCs were also measured for reference. For this, an LN cell suspension was positively selected for  $CD11c^{+}$  cells and stained with various combinations of  $CD11c$ , Langerin (intracellular), and  $CD8$ . Naive  $CD44$ -deficient and wild-type mice showed comparable numbers of Langerin $^{+}$  DCs,  $CD8^{+}$  DCs, and total  $CD11c^{+}$  DCs in their LNs (Figure 1d). Next, we examined the number of Langerin $^{+}$  DCs at 4 days after epicutaneous sensitization with protein antigen, which was the peak time point when epidermal LCs arrived



**Figure 2. T-cell responses in CD44-deficient and wild-type C57BL/6 mice.** Groups of CD44-deficient or wild-type mice were immunized by (a) patch application with ovalbumin (OVA) solution or phosphate-buffered saline control for 5 successive days or (b) subcutaneous injection with OVA plus complete Freund's adjuvant on day 1. At 10 days after immunization, draining lymph nodes (LNs) were isolated and *in vitro* OVA reactivation cultures were done. IFN- $\gamma$ , IL-5, and IL-13 levels in 48-hour supernatants were determined by ELISA. Net concentration (concentration in the absence of OVA subtracted from the concentration in the presence of OVA) is shown. Representative results from three independent experiments are shown.

in draining LNs. As shown in Figure 1e, CD44-deficient mice had higher numbers of CD11c<sup>+</sup> DCs in draining LNs compared with wild-type mice. Wilcoxon's rank sum test was used to test the significance of difference. The *P*-value for CD11c<sup>+</sup> DC was 0.0079. The numbers of Langerin<sup>+</sup> DCs in CD44-deficient mice also seemed to be higher than those in wild-type mice. However, it did not reach statistical significance. To confirm that these Langerin<sup>+</sup> DCs were skin-derived, fluorescence-conjugated ovalbumin (OVA-DQ) was used. Figure 1f shows that Langerin<sup>+</sup> DCs in draining LNs carried this fluorescent marker. We also checked the expressions of costimulating molecules on Langerin<sup>+</sup> DCs in draining LNs at 96 hours after epicutaneous sensitization with protein antigen. Langerin<sup>+</sup> DCs in draining LNs in CD44-deficient mice showed comparable expressions of CD80, CD86, CD24, and CD40 with those in wild-type mice (data not shown). Recently, it was recognized that epidermal Langerin<sup>+</sup> DCs (LCs) and dermal Langerin<sup>+</sup> DCs could be distinguished on the basis of their expressions of CD103 (Merad *et al.*, 2008). Figure 1g shows that the numbers of CD103<sup>+</sup> Langerin<sup>high</sup> DCs, which indicated epidermal LCs in draining LNs 96 hours after epicutaneous immunization, were comparable in CD44-deficient

and wild-type mice. The discrepancy between the results of Langerin<sup>+</sup> DCs and CD103<sup>+</sup> Langerin<sup>high</sup> DCs could be because of variations in CD103<sup>high</sup> Langerin<sup>high</sup> dermal DCs and CD103<sup>low</sup> CD8<sup>+</sup> Langerin<sup>low</sup> resident DCs. Taken together, these results show that CD44-deficient mice do not exhibit impairment of epidermal LC migration to LNs after epicutaneous sensitization with protein antigen.

We next checked T-cell immune responses in CD44-deficient mice using two prototypic experimental systems. Epicutaneous sensitization with protein antigen induces a predominant T helper (Th)2 response, whereas subcutaneous immunization induces a vigorous Th1 response. At 10 days after the start of immunization, drained LNs were isolated and used for *in vitro* antigen reactivation cultures. The interleukin-5 and interleukin-13 levels in supernatants of CD44-deficient mice with ovalbumin patch applications were always higher compared with those of wild-type mice (Figure 2a). However, the interferon- $\gamma$  levels in supernatants of CD44-deficient mice after subcutaneous immunization with ovalbumin plus complete Freund's adjuvant (CFA) were also higher than those from wild-type mice (Figure 2b). Thus, T-cell immune responses of CD44-deficient mice are not Th1 or Th2 biased. In

contrast, CD44-deficient mice showed stronger predominant Th responses after immunization.

Recently, CD44 was found to be a phagocytic receptor and involved in the resolution of noninfectious tissue inflammation by mediating the clearance of fragmented hyaluronan (Teder *et al.*, 2002; Vachon *et al.*, 2006). However, it is unlikely that these characteristics contributed to our results. First, we used gentle shaving to avoid harming the skin and applied patches without previous tape stripping or adding adjuvant. Thus, the protein-patch model that we established one decade ago examined epicutaneous sensitization under physiological, noninflammatory cutaneous conditions. Second, because fragmented hyaluronan can activate DCs, our results showing comparable expressions of costimulatory molecules on Langerin<sup>+</sup> DCs support the notion that the patched skin of CD44-deficient mice was noninflammatory. Third, CD44-deficient mice exhibit unremitting inflammation 7 days after injury, which was beyond our immunization schedule.

In summary, by using specific markers and an optimal time point, this study unexpectedly showed that CD44-deficient mice do not exhibit impairment of epidermal LC migration to LNs after epicutaneous sensitization with protein antigen.

# CONFLICT OF INTEREST

The authors state no conflict of interest.

# ACKNOWLEDGMENTS

This work was supported by a grant from the National Health Research Institutes (NHRI-EX98-9516SC). This work was supported in part by the Department of Medical Research of NTUH.

Shi-Chuen Miaw<sup>1</sup>, Jau-Shiuh Chen<sup>2</sup>,  
Pei-Chen Hsieh<sup>2</sup>, Ching-Yi Liu<sup>2</sup>,  
Ming-Hui Chung<sup>2</sup>, Pei-Chun Chen<sup>2</sup>,  
Mei-Ling Chang<sup>3</sup>, Fen-Ling Chen<sup>3</sup>,  
John T. Kung<sup>3</sup> and Li-Fang Wang<sup>2</sup>

<sup>1</sup>Graduate Institute of Immunology, National Taiwan University Hospital and National Taiwan University College of Medicine, Taipei, Taiwan; <sup>2</sup>Department of Dermatology, National Taiwan University Hospital and National Taiwan University College of

Medicine, Taipei, Taiwan and <sup>3</sup>Institute of Molecular Biology, Academia Sinica, Taipei, Taiwan  
E-mail: lifangwa@ntu.edu.tw or  
mbkung@ccvax.sinica.edu.tw

# REFERENCES

- Bursch LS, Wang L, Igyarto B *et al.* (2007) Identification of a novel population of Langerin+ dendritic cells. *J Exp Med* 204: 3147–56
- Chen JS, Chiu HC, Hsu CJ *et al.* (2009) Low-energy visible light irradiation modulates immune responses induced by epicutaneous sensitization with protein antigen. *J Invest Dermatol* 129:2258–64
- Isacke CM (2002) Molecules in focus the hyaluronan receptor, CD44. *Int J Biochem Cell Biol* 34:718–21
- Merad M, Ginhoux F, Collin M (2008) Origin, homeostasis and function of Langerhans cells

and other langerin-expressing dendritic cells. *Nat Rev Immunol* 8:935–47

- Mummert DI, Takashima A, Mummert ME (2004) Langerhans cells in CD44-deficient mice emigrate from the epidermis but fail to reach the lymph nodes after hapten application. *J Invest Dermatol* 122:846–7
- Shklovskaya E, Roediger B, Fazekas de St Groth B (2008) Epidermal and dermal dendritic cells display differential activation and migratory behavior while sharing the ability to stimulate CD4+ T cell proliferation *in vivo*. *J Immunol* 181:418–30
- Teder P, Vandivier RW, Jiang D *et al.* (2002) Resolution of lung inflammation by CD44. *Science* 296:155–8
- Vachon E, Martin R, Plumb J *et al.* (2006) CD44 is a phagocytic receptor. *Blood* 107:4149–58
- Weiss JM, Sleeman J, Renkl AC *et al.* (1997) An essential role for CD44 variant isoforms in epidermal Langerhans cell and blood dendritic cell function. *J Cell Biol* 137:1137–47


See related commentary on pg 2535

# Hair Follicles from Alopecia Areata Patients Exhibit Alterations in Immune Privilege-Associated Gene Expression in Advance of Hair Loss

*Journal of Investigative Dermatology* (2010) **130**, 2677–2680; doi:10.1038/jid.2010.180; published online 8 July 2010

# TO THE EDITOR

Alopecia areata (AA) is a putative autoimmune hair loss disease. Anagen-stage hair follicles may have immune privilege (Billingham and Silvers, 1971) comprising downregulation of major histocompatibility complex (MHC) class I and II and production of immunoregulatory agents (Paus *et al.*, 2003, 2005; Ito *et al.*, 2004). Of several hypotheses for AA pathogenesis (McElwee *et al.*, 2002; Gilhar *et al.*, 2007), one suggests that AA is a consequence of immune privilege collapse (Paus *et al.*, 2003; Ito *et al.*, 2004). Although immunohistological studies on immune privilege in AA have been performed (McDonagh *et al.*, 1993; Ito *et al.*, 2004), expression has not been objectively quantified and some immune privilege factors have not been investigated. We conducted



**Figure 1. Differential expression of immune privilege-associated genes in the lesional margin of alopecia areata (AA)-affected patches and the perilesional scalp hair follicles as compared with normal scalp specimens.** Quantitative PCR (qPCR) revealed apparent marked increased expression of IL1 receptor antagonist (IL1Ra), CD80, and macrophage inhibitory factor (MIF) in lesional margin hair follicle bulbs. Red/IK cytokine was significantly downregulated in the lesional margin. The fold change of gene expression was analyzed by qPCR and was calculated using the formula  $2^{-\Delta\Delta Ct}$ , using perilesional and lesional hair follicles from four AA patients versus normal hair follicles from four controls with  $\geq 20$  hair follicles per biopsy, and qPCR analysis was conducted in triplicate. Error bars represent the range factor difference ( $2^{-\Delta\Delta Ct \pm \Delta Ct}$  standard deviation). \*Indicates  $P < 0.05$  by Student's *t*-test. The y axis is in log10 scale.